



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			

Display	Abstract	▼	Sort	▼	Save	Text	Add to Clipboard	Order
---------	----------	---	------	---	------	------	------------------	-------

1: Environ Mol Mutagen 2000;36(1):5-12

[Related Articles](#), [Books](#), [LinkC](#)



## **DNA degradation by the mixture of copper and catechol is caused by DNA-copper-hydroperoxo complexes, probably DNA-Cu(I)OOH.**

**Schweigert N, Acero JL, von Gunten U, Canonica S, Zehnder AJ, Eggen RI.**

Swiss Federal Institute for Environmental Science and Technology (EAWAG),  
Dübendorf, Switzerland.

Free hydroxyl radicals (free ( $\cdot$ )OH), singlet oxygen ( $(^1\text{O}_2)$ ), or ( $\cdot$ )OH produced by DNA-copper-hydroperoxo complexes are possible DNA-damaging reactive oxygen species (ROS) in the reaction system containing copper, catechol, and DNA. para-Chlorobenzoic acid (pCBA) degradation studies revealed that  $\text{CuCl}_2$  mixed with catechol produced free ( $\cdot$ )OH. In the presence of DNA, however, inhibition of the pCBA degradation suggested that another ROS is responsible for the DNA degradation. Of a series of ROS scavengers investigated, only KI, NaN<sub>3</sub>, and Na-formate—all of the salts tested—strongly inhibited the DNA degradation, suggesting that the ionic strength rather than the reactivity of the individual scavengers could be responsible for the observed inhibition. The ionic strength effect was confirmed by increasing the concentration of phosphate buffer, which is a poor ( $\cdot$ )OH scavenger, and was interpreted as the result of destabilization of DNA-copper-hydroperoxo complexes. Piperidine-labile site patterns in DNA degraded by copper and catechol showed that the mixture of Cu(II) and catechol degrades DNA via the intermediate formation of a DNA-copper-hydroperoxo complex. Replacement of guanine by 7-deazaguanine did not retard the DNA degradation, suggesting that the DNA-copper-hydroperoxo complexes do not bind to the guanine N-7 as proposed in the literature. Copyright 2000 Wiley-Liss, Inc.

PMID: 10918354 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Add to Clipboard	Order
---------	----------	---	------	---	------	------	------------------	-------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)